



Junior Cert Maths

Free Notes

Quadratic Equations



Quadratic Equations

Quadratic equations are equations of the form $ax^2 + bx + c = 0$.

To solve a quadratic equation we must firstly make sure our equation equals zero. For instance if $x^2 + 8x + 8 = -7$ we must bring -7 to the other side of our equation so as it equals 0 i.e. $x^2 + 8x + 15 = 0$

Once our equation equals 0 it can be solved by factorisation. We let both of our factorised terms equal 0 and solve for x

$$\begin{aligned}x^2 + x - 42 &= 0 \\(x - 6)(x + 7) &= 0 \\x - 6 = 0 \quad x + 7 = 0 \\x = 6 \quad x = -7\end{aligned}$$

$$\begin{aligned}3x^2 + 19x + 26 &= 0 \\(3x + 13)(x + 2) &= 0 \\3x + 13 = 0 \quad x + 2 = 0 \\3x = -13 \quad x = -2 \\x = -13/3\end{aligned}$$

$$\begin{aligned}x^2 + 7x - 44 &= 0 \\(x + 11)(x - 4) &= 0 \\x + 11 = 0 \quad x - 4 = 0 \\x = -11 \quad x = 4\end{aligned}$$

$$\begin{aligned}6x^2 - 29x + 30 &= 0 \\(3x - 10)(2x - 3) &= 0 \\3x - 10 = 0 \quad 2x - 3 = 0 \\3x = 10 \quad 2x = 3 \\x = 10/3 \quad x = 3/2\end{aligned}$$

$$\begin{aligned}2x^2 - 15x + 27 &= 0 \\(2x - 9)(x - 3) &= 0 \\2x - 9 = 0 \quad x - 3 = 0 \\2x = 9 \quad x = 3 \\x = 9/2\end{aligned}$$

$$\begin{aligned}x^2 - 6x &= 0 \\x(x - 6) &= 0 \\x = 0 \quad x - 6 = 0 \\x = 0 \quad x = 6\end{aligned}$$

$$\begin{aligned}12x^2 - 7x - 45 &= 0 \\(4x - 9)(3x + 5) &= 0 \\4x - 9 = 0 \quad 3x + 5 = 0 \\4x = 9 \quad 3x = -5 \\x = 9/4 \quad x = -5/3\end{aligned}$$

$$\begin{aligned}2x^2 + 18x &= 0 \\2x(x + 9) &= 0 \\2x = 0 \quad x + 9 = 0 \\x = 0 \quad x = -9\end{aligned}$$

If a quadratic equation cannot be factorised we must use the quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

where a, b, c are taken from the quadratic equation $ax^2 + bx + c = 0$
Use your calculator to help you solve these types of quadratic equations.

Solve $x^2 + 5x + 3 = 0$

a = 1
b = 5
c = 3

$$\frac{-5 \pm \sqrt{5^2 - 4(1)(3)}}{2(1)}$$

$$\frac{-5 \pm \sqrt{25 - 12}}{2}$$

$$\frac{-5 \pm \sqrt{13}}{2}$$

We Split our equation in two

$$\frac{-5 + \sqrt{13}}{2} \quad \frac{-5 - \sqrt{13}}{2}$$

Using our calculator we get

x = -0.7 and x = -4.3

Questions

1. Solve for x: $3x^2 + 11x = 4$

$$3x^2 + 11x = 4$$

$$3x^2 + 11x - 4 = 0$$

Solve $2x^2 - 4x - 3 = 0$

a = 2
b = -4
c = -3

$$\frac{4 \pm \sqrt{(-4)^2 - 4(2)(-3)}}{2(2)}$$

$$\frac{4 \pm \sqrt{16 + 24}}{4}$$

$$\frac{4 \pm \sqrt{40}}{4}$$

We Split our equation in two

$$\frac{4 + \sqrt{40}}{4} \quad \frac{4 - \sqrt{40}}{4}$$

Using our calculator we get

x = -2.58 and x = -0.58

$$(3x - 1)(x + 4)$$

$$3x - 1 = 0 \quad x + 4 = 0$$

$$3x = 1 \quad x = -4$$

$$x = 1/3$$

$$x = 1/3 \text{ and } -4$$

2. Solve for x $8x^2 - 14x + 3 = 0$

$$8x^2 - 14x + 3 = 0$$

$$(2x - 3)(4x - 1)$$

$$2x - 3 = 0 \quad 4x - 1 = 0$$

$$2x = 3 \quad 4x = 1$$

$$x = 3/2 \quad x = 1/4$$

$$x = 3/2 \text{ and } 1/4$$

3. Find the roots of the equation $2x^2 - 7x - 6 = 0$ Give your answers correct to two decimal places

Use the quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$a = 2$$

$$b = -7$$

$$c = -6$$

$$\frac{-7 \pm \sqrt{(-7)^2 - 4(2)(-6)}}{2(2)}$$

$$\frac{-7 \pm \sqrt{49 + 48}}{4}$$

$$\frac{-7 \pm \sqrt{97}}{4}$$

Split our equation into two parts

$$\frac{-7 + \sqrt{97}}{4} \quad \frac{-7 - \sqrt{97}}{4}$$

Using our calculator we get

$$x = 4.21 \text{ and } -0.71$$

4(i). Solve the equation $x^2 = 3x + 2$. Give your answers correct to two decimal Places

$$x^2 - 3x - 2 = 0$$

Use the quadratic formula

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$a = 1$$

$$b = -3$$

$$c = -2$$

$$\frac{3 \pm \sqrt{(-3)^2 - 4(1)(-2)}}{2(1)}$$

$$\frac{3 \pm \sqrt{9+8}}{2}$$

$$\frac{3 \pm \sqrt{17}}{2}$$

Split our equation into two parts

$$\frac{3+\sqrt{17}}{4} \quad \frac{3-\sqrt{17}}{4}$$

Using our calculator we get

$$x = 3.56 \text{ and } -0.56$$

**4(ii) Hence, or otherwise, find value for p for which $p = 3\sqrt{p} + 2$
Give your answers correct to one decimal place.**

From our previous equation $x^2 = 3x + 2$ $x = 3.56$ and -0.56

If we replace x with \sqrt{p}

$$\sqrt{p} = 3.56 \quad \sqrt{p} = -0.56$$

$$p = 12.7 \quad p = 0.3$$

If we test both of these values back into our equation $p = 3\sqrt{p} + 2$

we find that only 12.7 satisfies our equation
 $p = 12.7$

5. (i) Solve the equation $x^2 - 6x + 4 = 0$, giving your answer in the form of $a \pm b$, where a, b $\in \mathbb{N}$

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$a = 1$$

$$b = -6$$

$$c = 4$$

$$\frac{6 \pm \sqrt{(-6)^2 - 4(1)(4)}}{2(1)}$$

$$\frac{6 \pm \sqrt{36 - 16}}{2}$$

$$\frac{6 \pm \sqrt{20}}{2}$$

$$\frac{6 \pm 2\sqrt{5}}{2}$$

Split our equation into two parts

$$\frac{6 + 2\sqrt{5}}{2} \quad \frac{6 - 2\sqrt{5}}{2}$$

Dividing both equations across by 2 we get

$$x = 3 + \sqrt{5} \quad x = 3 - \sqrt{5}$$

(ii) Hence, or otherwise, find two values for p for which $(3 + p)^2 - 6(3 + p) + 4 = 0$

From our previous equation $x^2 - 3x + 4 = 0$ $x = 3 + \sqrt{5}$ and $x = 3 - \sqrt{5}$

If we replace x with $3 + p$ we get the given equation $(3 + p)^2 - 6(3 + p) + 4 = 0$

So $x = 3 + p$

$$p + 3 = 3 + \sqrt{5} \quad \text{and} \quad p + 3 = 3 - \sqrt{5}$$

$$p = \sqrt{5} \quad \text{and} \quad p = -\sqrt{5}$$

(iii) Show that the sum of the two values of p is zero.

$$\sqrt{5} - \sqrt{5} = 0$$

For more comprehensive Junior Cert Revision Notes Click
Here.... [Junior Cert Maths Notes](#)