



Differentiation
Maths Past Exam Questions
Higher Level

2012

Paper 1 – Section A – Q6

6. (a) Differentiate with respect to x :

(i) $(4x^2 - 1)^3$.

(ii) $\sin^{-1}\left(\frac{2x}{3}\right)$.

(b) (i) Differentiate \sqrt{x} with respect to x , from first principles.

(ii) Find the equation of the tangent to the curve $y = \sqrt{x}$ at the point $(9, 3)$.

(c) Let f be the function $f : x \rightarrow 8x + \sin 4x + 4 \sin 2x$, where $x \in \mathbb{R}$.

(i) Find $f'(x)$.

(ii) Express $f'(x)$ in terms of $\cos 2x$.

(iii) Prove that $f(x)$ is increasing for all values of x .

Paper 1 – Section A – Q6

6. (a) Differentiate $\cos^2 x$ with respect to x .
- (b) The equation of a curve is $y = e^{-x^2}$.
- Find $\frac{dy}{dx}$.
 - Find the co-ordinates of the turning point of the curve.
 - Determine whether this turning point is a local maximum or a local minimum.
- (c) The function f is defined as $x \rightarrow \frac{2x}{x+1}$, where $x \in \mathbb{R} \setminus \{-1\}$.
- Find the equations of the asymptotes of the curve $y = f(x)$.
 - P and Q are distinct points on the curve $y = f(x)$.
The tangent at Q is parallel to the tangent at P .
The co-ordinates of P are $(1, 1)$.
Find the co-ordinates of Q .
 - Verify that the point of intersection of the asymptotes is the midpoint of $[PQ]$.

2010

Paper 1 – Section A – Q6

6. (a) The equation $x^3 + x^2 - 4 = 0$ has only one real root.
Taking $x_1 = \frac{3}{2}$ as the first approximation to the root, use the Newton-Raphson method to find x_2 , the second approximation.

- (b) Parametric equations of a curve are:

$$x = \frac{2t-1}{t+2}, \quad y = \frac{t}{t+2}, \quad \text{where } t \in \mathbb{R} \setminus \{-2\}.$$

(i) Find $\frac{dy}{dx}$.

- (ii) What does your answer to part (i) tell you about the shape of the graph?

- (c) A curve is defined by the equation $x^2y^3 + 4x + 2y = 12$.

(i) Find $\frac{dy}{dx}$ in terms of x and y .

- (ii) Show that the tangent to the curve at the point $(0, 6)$ is also the tangent to it at the point $(3, 0)$.

2009

Paper 1 – Section A – Q6

6. (a) Differentiate $\sin(3x^2 - x)$ with respect to x .
- (b) (i) Differentiate \sqrt{x} with respect to x , from first principles.
- (ii) An object moves in a straight line such that its distance from a fixed point is given by $s = \sqrt{t^2 + 1}$, where s is in metres and t is in seconds.
Find the speed of the object when $t = 5$ seconds.
- (c) The equation of a curve is $y = \frac{2}{x-3}$.
- (i) Write down the equations of the asymptotes and hence sketch the curve.
- (ii) Prove that no two tangents to the curve are perpendicular to each other.

2007

Paper 1 – Section A – Q6

6. (a) Differentiate $\frac{x^2 - 1}{x^2 + 1}$ with respect to x .
- (b) (i) Differentiate $\frac{1}{x}$ with respect to x from first principles.
- (ii) Find the equation of the tangent to $y = \frac{1}{x}$ at the point $\left(2, \frac{1}{2}\right)$.
- (c) Let $f(x) = \tan^{-1} \frac{x}{2}$ and $g(x) = \tan^{-1} \frac{2}{x}$, for $x > 0$.
- (i) Find $f'(x)$ and $g'(x)$.
- (ii) Hence, show that $f(x) + g(x)$ is constant.
- (iii) Find the value of $f(x) + g(x)$.

2006

Paper 1 – Section A – Q6

6. (a) Differentiate $\sqrt{x}(x+2)$ with respect to x .
- (b) The equation of a curve is $y = 3x^4 - 2x^3 - 9x^2 + 8$.
- (i) Show that the curve has a local maximum at the point $(0, 8)$.
- (ii) Find the coordinates of the two local minimum points on the curve.
- (iii) Draw a sketch of the curve.
- (c) Prove by induction that $\frac{d}{dx}(x^n) = nx^{n-1}$, $n \geq 1$, $n \in \mathbf{N}$.

Paper 1 – Section A – Q6

6. (a) Differentiate with respect to x

(i) $(1 + 7x)^3$ (ii) $\sin^{-1}\left(\frac{x}{5}\right)$.

(b) Let $y = \frac{1 - \cos x}{1 + \cos x}$.

Show that $\frac{dy}{dx} = t + t^3$, where $t = \tan \frac{x}{2}$.

(c) The equation of a curve is $y = \frac{x}{x-1}$, where $x \neq 1$.

- (i) Show that the curve has no local maximum or local minimum point.
- (ii) Write down the equations of the asymptotes and hence sketch the curve.
- (iii) Show that the curve is its own image under the symmetry in the point of intersection of the asymptotes.